

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 17:24:46 ON 19 MAR 2006

=> fil pctfull

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

0.21

0.21

FILE 'PCTFULL' ENTERED AT 17:24:57 ON 19 MAR 2006

COPYRIGHT (C) 2006 Univentio

FILE LAST UPDATED:

14 MAR 2006

<20060314/UPTX>

MOST RECENT UPDATE WEEK:

200610

FILE COVERS 1978 TO DATE

>>> IMAGES ARE AVAILABLE ONLINE AND FOR EMAIL-PRINTS <<<

>>> NEW IPC8 DATA AND FUNCTIONALITY NOT YET AVAILABLE IN THIS FILE.
USE IPC7 FORMAT FOR SEARCHING THE IPC. WATCH THIS SPACE FOR FURTHER
DEVELOPMENTS AND SEE OUR NEWS SECTION FOR FURTHER INFORMATION

>>> UPDATING OF BIBLIOGRAPHIC DATA DELAYED DUE TO DELIVERY
FORMAT CHANGES <<<

>>> FULL-TEXT UPDATES CONTINUE. PATENT NUMBER AVAILABLE FOR DISPLAY
ONLY, USE FIELD CODE FPI <<<

>>> SDI SEARCHES (ALERTS) WILL BE RESUMED WHEN BIBLIOGRAPHIC DATA
BECOME AVAILABLE <<<

=> WO96/20236/pn

L1 0 WO96/20236/PN
(WO96/PN)

=> WO 1996020236/pn

L2 1 WO 1996020236/PN
(WO9620236/PN)

=> form? and l2

952912 FORM?
L3 1 FORM? AND L2

=> d kwic l3

L3 ANSWER 1 OF 1 PCTFULL COPYRIGHT 2006 Univentio on STN

PI WO 9620236 A1 19960704

ABEN A polyglycomer is disclosed of **formula** (I), wherein L is a
linker selected from the group of
-O-(CH₂)_m-, -NH-(CH₂)_m-, -O-(CH₂)_m-X-, and -NH-(CH₂)_m-X-, wherein X = S.

ABFR L'invention se rapporte a un polyglycomere de la **formule** (I)
dans laquelle L represente un
element de liaison selectionne dans le groupe de -O- (CH₂)_m-,
-NH-(CH₂)_m-, -O-(CH₂)_m-X-, et
-NH-(CH₂)_m-X-, . . .

DETD . . . the Invention

In general, the field of the present invention is
carbohydrate-substituted polymers. Specifically, the field of
the present invention is saccharide-substituted polymers
5 **formed** from 7-oxanorbornenes through aqueous ring-opening

metathesis polymerization.

Summary of the Invention

The present invention is a compound of the **formula**.

The present invention is also a polyglycomer of the **formula**.

0

0

L L

\ 2

R' R

wherein L is a linker selected from the group of (CH₂),-, -NH-(CH₂)_m-f (CH₂),-X- and -NH-(CH₂),-X- wherein X. . . and m is 2 - 10; and wherein R' and R 2 are selected from the group consisting of moieties of the **formula**.

The present invention is also a compound of the **formula**

0

L 0

/ CH₃

R

5 wherein L is a linker selected from the group of (CH₂),-f NH-(CH₂)_m-f (CH₂)_m-X- and -NH-(CH₂),-X- moieties, wherein X 0 or. . .

The present invention is also a compound of the **formula**

0

Hoes etlH

0 0

L L

R' R₂

wherein L is a linker selected from the group of (CH₂),-, NH-(CH₂)_m-f (CH₂),-X- and -NH-(CH₂)_m-X- moietiesf wherein X 0. . .

via a C-glycoside linkage and treating the product of the first step with ruthenium trichloride, wherein a polyglycomer is created. In a preferred **form** of this method, the ruthenium trichloride treatment is in a aqueous environment at 550C.

When the saccharide is attached with a C-glycoside linker, the linker element of the general **formula** described 2 5 above is CH₂-CH₂-'. However, we envision that other linkers WO 96/20236 PCrFUS95/13361

9

hexoses and pentoses are suitable, although hexoses. . .

designs of 8, 9 and 10 are based on the Lewis a rather than Lewis x template, because derivatives of the **former** generally bind more tightly to the selectins (Berg, et al., Biochem. Biophys. Res. Commun. 184:1048-1055, 1992. Nelson, et al., J. Clin. Invest. 91:1157-1166,. . .

selectin inhibitors, sulfates 8, 9 and 10, were generated by chemical synthesis. We chose to employ chemical over enzymatic synthesis because the **former** provides access to many different carbohydrate derivatives. Although enzymes are available for the generation of Le' structures, the corresponding Le' scaffold can not. . .

and the galactose partners 13a and 13b are available through standard manipulations (Manning, et al., 1994, in preparation, supplemental material). The glycosylations to **form** lactose derivatives 14a and 14b were promoted with Schmidt glycosylation conditions (Schmidt, In Comprehensive Organic Synthesis; B. M. Trost and I. Fleming, Eds.;. . .

Compound 8 was sulfated by way of the cyclic tin acetal, which **forms** at the 3 and 4 hydroxyls of the galactose residue.

CLMEN 1. A polyglycomer of the **formula**

0
L L-
R1 \ jt]
wherein L is a linker selected from the group of (CH2)m-f -
NH-(CH2)m-r (CH2)m-X-. and -NH-(CH2)m-X-j, wherein X is. . .

12 A polyglycomer of the **formula**

0
0 0
L L
RI
R2
wherein L is a linker selected from the group of (CH2),-, -
NH-(CH2)m-f (CH2)m-Xi and -NH-(CH2).-X-, wherein X is S or O
and m is 2-10,
wherein R' and R 2are selected from the group consisting of
moieties of the **formula**

HO
CH
H3C CH
HO 0 CH
R92o.0 0 0
R9b L
OH HO
wherein R'' = SO3Na and R 2 = H; or R SO3Na and R, . . .

13 A polyglycomer of the **formula**

n
0
RI CH3
wherein L is a linker selected from the group of (CH2),-, -
NH-(CH2)m-j, (CH2) M.X-1 and -NH-(CH2),,, -X-, wherein X is S. . .

14 A polyglycomer of the **formula**

0
His 41811H
0 0
L L
RI \ 2
R
wherein L is a linker selected from the group of (CH2),-, -
NH-(CH2)m-f (CH2),-X-1 and -NH-(CH2),-X-, wherein. . .

=> logoff hold
COST IN U.S. DOLLARS
FULL ESTIMATED COST

SINCE FILE	TOTAL
ENTRY	SESSION
14.73	14.94

SESSION WILL BE HELD FOR 60 MINUTES
STN INTERNATIONAL SESSION SUSPENDED AT 17:32:26 ON 19 MAR 200